




Ponderosa pine forests



Locoweed

©2004 Zackery Zdinak




Yarrow

© Robert DeWitt Ivey



Mullein

© Robert DeWitt Ivey




Fleabane

© Robert DeWitt Ivey



Rocky Mountain iris

©1999 Zackery Zdinak




Bracken fern

©1999 Zackery Zdinak



Silvery lupine

©1999 Zackery Zdinak




Purple geranium

© Robert DeWitt Ivey




Scarlet gilia

©1999 Zackery Zdinak



Whitemargin pussytoes*



Pingue rubberweed

© Robert DeWitt Ivey



Mountain parsley

© Robert DeWitt Ivey

Locoweed (*Oxytropis lambertii*)
• $\frac{3}{4}$ " long purple pea-like flowers in clusters to 9" tall. Leaves to 5" long with 11–17 fuzzy leaflets, sometimes appearing gray. Seedstalk turns pale yellow and stays standing after releasing seed.

Rocky Mountain iris, blue flag (*Iris missouriensis*) • Common in moist sites, and into the mixed conifer zone. Showy purplish petals and white sepals with blue stripes on round stem to 24" tall. Long, narrow leaves from base of plant. Grows in groups. Growth depends on adequate snowfall.

Scarlet gilia, skyrocket (*Ipomopsis aggregata*) • Bright red, trumpet-shaped flowers along stalk to 3' tall. Dark green leaves with narrow lobes. First year it consists of leaves close to the ground, second year it grows higher and then flowers.

Yarrow (*Achillea millefolium*) • Feathery leaves 2–6" long, $\frac{1}{2}$ " wide in rosettes of many leaves. White flat-topped clusters of flowers on stalk 6–20" tall.

Bracken fern (*Pteridium aquilinum*) • A large fern, to 3' tall. Fronds triangular, usually with three leaflets with many tiny leaves. Grows in large colonies in fields, brushy areas, and woods. Thick root-like underground stems, to 6' long; many fronds may grow from each stem. Vigorous after disturbance. Cover for small animals and emerging plants.

Whitemargin pussytoes (*Antennaria marginata*) • Small plant lying close to ground forming mats. Necktie shaped leaves $\frac{1}{2}$ –1" long, slightly hairy underneath, with a white appearance. White clusters of flowers on stalks 2–4" tall.

Mullein (*Verbascum thapsus*) • Readily invades disturbed sites but only persists for a few years. First year forms basal rosette of large soft, fuzzy leaves, second year sends up seed stalk to 7' high. Hundreds of bright yellow flowers maturing to round-seed pods of tiny black seeds. Seed stalk dries to a stout brown shaft that persists through winter.

Silvery lupine (*Lupinus argenteus*) • Branching flowers to 3' tall. Leaves fuzzy when young; 5–9 narrow lobes; pea-like flowers in tall clusters; purple. Aggressive growth after fire.

Pingue rubberweed (*Hymenoxys richardsonii*) • To 12" tall, several to dozens of woody stems in a bunch, wooly at base. Yellow daisy-like flowers $\frac{1}{2}$ " wide maturing to wooly tufts of seeds. Very narrow leaves, poisonous to livestock.

Fleabane (*Erigeron* spp.) • Many species. Small white, pink, or light purple flowers with numerous narrow petals and yellow center.

Purple geranium (*Geranium caespitosum*) • Pink or purple flowers; up to 10" tall; few flowers per plant. Thin creeping stems, often red. Leaves five-lobed, smelly when crushed, red in autumn.

Mountain parsley (*Pseudocymopterus montanus*) • Up to 30" tall. Yellow flowers in flat-topped clusters. Leaves green, broad, feathery, with many pointed lobes. Strong scent.

* Cronquist, A. and A. H. Holmgren, N. H. Holmgren, J. L. Reveal and P. K. Holmgren. 1994. *Intermountain Flora Vascular Plants of the Intermountain West, USA*. The New York Botanical Garden, Bronx, NY. Volume 5.

Wildflowers

